



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

APRIL 16.

The President, Dr. JOSEPH LEIDY, in the chair.

Thirty-four persons present.

The following papers were presented for publication :—

“A contribution to the Spider fauna of the Bermuda Islands,”
by Dr. Geo. Marx.

“Notes on a small collection of Myriapods from the Bermuda Islands,” by Charles H. Bollman.

“A contribution to the Actinology of the Bermudas,” by J. Playfair McMurrich, Ph. D.

“On some new species of Mollusca from the Bermuda Islands,”
by Angelo Heilprin.

A Parasitic Copepod.—Prof. LEIDY remarked that last summer while at Beach Haven, N. J., there was brought to him from the surf a living specimen of the singular, transparent fish *Leptocephalus*. In examining it he observed attached to the tail-fin a minute copepod crustacean, apparently of the genus *Chalimus*. The parasite was attached by a long filiform rostrum and resembled in this and other respects more the *Chalimus scomberi*, as represented by Baird, in fig. 5, tab. xxxiii, of the British Entomostraca, than it does the original of this species as represented by Burmeister in the Nova Acta N. C. of Bonn, xvii, tab. 23, fig. 13. The species which may be distinguished as *Chalimus tenuis* is considerably less than half the size of *C. scomberi*. The cephalothorax nearly twice the length of the breadth, is obcordate and proportionately much narrower than in the latter species. The frontal segment is narrow and not prominent laterally, and the biarticulate antennae are concealed beneath. The abdomen, half the length of the cephalothorax, exhibits three conspicuous divisions, and the short caudal appendages end in three minute setae. Abdominal feet ending in biramous leaf-like segments fringed with short setae. Rostrum linear and almost as long as the cephalothorax. Whole length 1.125 mm; length of cephalothorax 0.5 mm; breadth 0.275; length of rostrum 10.5; length of abdomen 0.25 mm.



The accompanying outline represents the animal magnified forty-four diameters.